

Degenerative changes of the spine in people from prehistoric Okhotsk culture and two ancient human groups from Kanto and Okinawa, Japan

Yasushi SHIMODA¹, Tomohito NAGAOKA², Keiichi MOROMIZATO¹, Masanobu SUNAGAWA¹,
Tsunehiko HANIHARA³, Minoru YONEDA⁴, Kazuaki HIRATA², Hiroko ONO⁵,
Tetsuya AMANO⁵, Tadahiko FUKUMINE¹, Hajime ISHIDA^{1*}

¹*Department of Human Biology and Anatomy, Faculty of Medicine, University of the Ryukyus, Nishihara 903-0215, Japan*

²*Department of Anatomy, St. Marianna University School of Medicine, Kawasaki 216-8511, Japan*

³*Department of Anatomy, Kitasato University School of Medicine, Sagami 252-0374, Japan*

⁴*Department of Integrated Biosciences, Graduate School of Frontier Sciences, The University of Tokyo, Kashiwa 277-8562, Japan*

⁵*Hokkaido University Museum, Hokkaido University, Sapporo 060-0810, Japan*

Received 25 September 2010; accepted 11 April 2011

Abstract Degenerative changes of the spine in people of the Okhotsk culture were investigated in adult human skeletal remains from 38 males and 34 females. These findings were then compared with those in materials obtained from the medieval Kamakura period and early-modern peasants on Kumejima, Ryukyu Islands. The three samples clearly showed different patterns. In the Okhotsk series, the cervical spine of each sex had most osteophytes on the vertebral body, while the Kumejima samples had the highest frequency on the lumbar vertebrae. In the Kamakura series, males were most affected on the lower thoracic vertebrae. Moreover, severe osteophytes on the body of the lumbar vertebrae were more frequently seen in the Okhotsk males. Degenerative changes of the articular process of the Okhotsk series were most frequently seen in the lumbar vertebrae and least frequently seen in the cervical vertebrae. This is well contrasted with a high frequency of degenerative changes of cervical apophyseal joint in early-modern Kumejima peasants. The Kamakura series of each sex had generally low frequencies. Severe degenerative changes of apophyseal joint dominantly affected the Okhotsk series. It is inferred that different dynamic loads caused a high frequency of degenerative changes in the corresponding articular parts. For example, because the Okhotsk culture developed a considerable maritime infrastructure, the lifestyle required for sea-mammal hunting and fishing seems to have particularly affected the incidence of severe degenerative changes of the lumbar vertebrae.

Key words: degenerative changes, osteoarthritis, spine, Okhotsk culture, paleopathology

Introduction

Degenerative joint disease, or osteoarthritis, is a common disease characterized pathologically by damage to the articular cartilage and bone surface in synovial joints, associated with osteophyte formation at the joint margins (Dieppe and Lohmander, 2005). This condition affects not only the hands, hips and knees, but also the spine, and is strongly age-related, currently rising in frequency after 50 years old (Felson et al., 2000). In addition to aging and systematic factors such as genetics, obesity and sex, the other risk factors for osteoarthritis consist of joint injuries and specific repetitious activities, or chronic overload. For example, with regard to occupational factors, farmers tend to have osteoarthritis of hip joints and workers undertaking physical

labor have high frequencies of knee osteoarthritis (Coggon et al., 1998; Felson et al., 2000). Although joint damage and clinical joint pain are both common, it is clear that the severity of the joint damage on a radiograph bears little relation to clinical severity (Dieppe and Lohmander, 2005).

Degenerative change of the spine is a common age-related vertebral alteration (Gallucci et al., 2005; Roh et al., 2005; Freund and Sartor, 2006; Pytel et al., 2006; Kalichman and Hunter, 2007; Ruan et al., 2007; Shedid and Benzel, 2007). The vertebral bodies are interconnected by the intervertebral discs (fibrocartilage), which are called symphyses. On the other hand, the apophyseal joints between articular processes in the vertebral arches are normally synovial joints. The typical osteological findings consist of anterolateral osteophytes where Sharpey's fibers attach to the vertebral body, degenerative change, or facet arthrosis of apophyseal joints (Resnick, 2002). Osteophytes on the vertebral body are more frequent in men than in women, as well as in older populations, and also appear to affect persons engaged in heavy physical labor. They may affect any segment of the vertebral column. Because degenerative changes of apophyseal joints are also common, some pathological evidence of such

* Correspondence to: Hajime Ishida, Department of Human Biology and Anatomy, Faculty of Medicine, University of the Ryukyus, Uehara 207, Nishihara, Okinawa 903-0215, Japan.
E-mail: ishidaha@med.u-ryukyu.ac.jp

Published online 25 June 2011

in J-STAGE (www.jstage.jst.go.jp) DOI: 10.1537/ase.100925

disease is seen in the spine of all individuals after the age of 50 or 60 years. Degenerative changes of apophyseal joints commonly affect the middle and lower cervical spine, upper and mid-thoracic spine, and the lower lumbar spine (Resnick, 2002).

Because risk factors for degenerative disease of the spine consist of not only age but also selected activities, many physical anthropologists have investigated degenerative changes of the spine from prehistoric and historic human populations in order to reconstruct their life activity (Suzuki, 1978, 1998; Fukushima, 1988; Bridges, 1994; Lovell, 1994; Lieverse et al., 2007; Moromizato et al., 2007; Rojas-Sepúlveda et al., 2008). In general, upright bipedal humans exhibit more marked degenerative changes of the spine than other hominoids (Jurmain, 2000). As mentioned above, although human spines tend to be more uniformly affected throughout the vertebral column, previous anthropological investigations revealed that prehistoric and historic human populations vary in terms of the prevalence of degenerative changes of the spine (Bridges, 1994; Moromizato et al., 2007). In addition, there are different patterns between the prevalence of osteophytes on the vertebral body and that of degenerative changes of apophyseal joint (Moromizato et al., 2007). This is one of the merits of studying degenerative changes of the spine in skeletal populations. However, because it is necessary to obtain large initial numbers of samples for epidemiological comparison, only a few studies have been performed in Japan (Suzuki, 1978, 1998; Fukushima, 1988).

In this study, we chose three ancient (historic) skeletal

collections of relatively large sample sizes with good preservation from the Japanese archipelago. The first is skeletal remains of the Okhotsk culture from Hokkaido and Sakhalin (Figure 1). The Okhotsk culture from the 5th to 12th century AD developed a considerable maritime infrastructure, which has been demonstrated by zooarcheological and isotopic analyses (Yoneda, 2002; Amano, 2003; Naito et al., 2010). The Okhotsk people concentrated on fishing and sea-mammal hunting. Many morphological and recent DNA analyses revealed that people from the Okhotsk culture were genetically similar to current Amur basin people (Ishida, 1996; Sato et al., 2007, 2009a, b; Komesu et al., 2008).

The second group of skeletal samples is from the medieval city of Kamakura, Kanto District, Central Japan (Figure 1). Medieval Kamakura was the capital of the Kamakura Shogunate in the period 1193–1333 AD. Huge numbers of human skeletal remains were recovered at the Yuigahama area, the coast of Kamakura, from the 1950s (Suzuki et al., 1956; Hirata et al., 2004). Because the samurai (military soldiers) were buried at a different place in Kamakura, the human remains from the Yuigahama area were considered to be those of common people. Mitochondrial DNA analysis revealed that the people of medieval Kamakura are very similar to the modern main-island Japanese (Shinoda, 2011).

Early-modern Kumejima samples from the Ryukyu Islands, the southernmost island chain of the Japanese archipelago, form the third group (Figure 1). One thousand human skeletal remains of the early-modern period were recovered from caves on Kumejima between 1998 and 2000




Figure 1. Location of the three skeletal series of the Okhotsk culture sites in Hokkaido, the Yuigahama-minami site of medieval Kamakura in Kanto, and the Yacchi-no-Gama site of the early-modern Kumejima, Okinawa, Japan.

(Fukumine et al., 2001, 2006; Irei et al., 2008). These caves had been used as a mortuary during the 17th–19th centuries AD. The past existence of paddy fields in this area indicates that the skeletal series represent farmers (Fukumine et al., 2001). Isotopic analysis indicated that these people consumed both C3 plants, which fix carbon via the Calvin–Benson cycle (e.g. taro, rice and tree), and fish (Yoneda et al., 2004; Irei et al., 2008).

Moromizato et al. (2007) reported on degenerative changes of the spine for the Kumejima samples in Japanese. They found that the lumbar regions were most affected for both sexes, while severe degenerative changes of the lumbar region were more frequently seen in females. In addition, it was suggested that the greater frequency of degenerative changes seen in female cervical vertebrae may have been caused by traditional behavior, such as carrying items on the head.

On the basis of this report on Kumejima (Moromizato et al., 2007), we would like to present basic data on degenerative changes of the spine among three different peoples from the Japanese archipelago, and to compare rates of prevalence among them. Finally, we will mention causes of the different prevalence patterns of degenerative changes among them on the basis of their subsistence, nutrition, labor, and culture.

Materials and Methods

Table 1 and Figure 1 show numbers and locations of the materials used in this study. In our analysis of sex-based and age-related differences in degenerative changes, we determined the age and sex of the individuals from which the skeletal materials were derived. Sex was estimated by standard methods, such as pelvic and cranial morphology (White, 2000). Age was also estimated by standard methods using auricular surfaces (Nagaoka et al., 2006), sutural closure, and dental attrition (White, 2000). Because the age distribution of the Kamakura people is different to those of the other peoples owing to their relatively short life expectancy (Nagaoka et al., 2006), we interpreted their prevalence rates carefully.

Although we used individuals of known sex and age with more than 20% vertebral preservation in order to obtain relative large sample sizes of respective vertebrae, the preservation rates were high. That is, the averages of preservation rates were 83% (female) and 92% (male) in the Kumejima

series, the averages were lower at 81% (female) and 85% (male) in the Kamakura series, and 62.9% (female) and 54.6% (male) in the Okhotsk series. No significant age-class differences were found.

For diagnosis of osteophytes on the vertebral body, the first author (Y.S.) examined osteophytes (bony outgrowth) on the body of vertebrae of the Okhotsk and Kamakura series, and the third author (K.M.) examined those of the Kumejima series (Moromizato et al., 2007). The osteophytes extend first horizontally and then in a vertical direction and finally bridge with the adjacent vertebrae (Resnick, 2002). We also examined progressive stages of osteophytes to evaluate severity (Figure 2). The normal situation is Grade 0; the next grade is when osteophytes grow horizontally (Grade 1); osteophytes can turn to grow in a vertical direction (Grade 2); osteophytes can then significantly grow in a vertical direction (Grade 3); eventually, osteophytes can bridge adjacent vertebrae (Grade 4), according to the diagnostic criteria of Rogers (1966) and Wada (1975). The grades were independently scored for the anterior, right and left sides, and posterior aspects of superior and inferior vertebral rims of the vertebral body. Differential diagnosis is very important but difficult because of lack of soft tissues, such as intervertebral discs. However, we could eliminate other bony outgrowths of the vertebral column that were clearly diagnosed as ankylosing spondylitis and psoriatic arthritis.

For diagnosis of degenerative changes of apophyseal joints, degenerative changes of the articular surface of the articular process were also examined by the first (Y.S.) and third (K.M.) authors. The pathological characteristics of osteoarthritis of the apophyseal joint are concordant with those seen in other synovial joints. Owing to a lack of articular cartilage, bony eburnation and sclerosis, pitting and osteophytes were checked. We also judged progressive stages of degenerative change of the apophyseal joint to evaluate severity (Figure 3). The normal condition is Grade 0; the next stage is when osteophytes grow on the rim of articular surface without pitting on the surface (Grade 1); osteophytes can also grow on the rim of articular surface with lipping with slight pitting (Grade 2); osteophytes can then grow all around the rim of the articular surface with moderate pitting on the surface and the rims of articular surface tend to be broken (Grade 3); eventually, osteophytes can significantly grow on the rim of articular surface with severe pitting on the surface and the rim becomes unclear (Grade 4),

Table 1. Materials used in this study

Sample	Date	Location	Sample size				Total	Institution	Source
			Female		Male				
			Younger (under 40)	Older (over 40)	Younger (under 40)	Older (over 40)			
Okhotsk	5th–12th AD	Hokkaido	14	20	23	15	72	HU, SMU	present study
Kamakura	13th AD	Kanto	42	30	50	22	144	STMU	present study
Kumejima	17th–18th AD	Ryukyu	16	29	21	35	101	OPAC	Moromizato et al. (2007)

HU: Hokkaido University, Sapporo, Japan.

SMU: Sapporo Medical University, Sapporo, Japan.

STMU: St. Marianna University, Kawasaki, Japan.

OPAC: Okinawa Prefectural Archaeological Center, Okinawa, Japan.




Figure 2. Progressive stages of osteophytes on the body of vertebrae. Grade 0: normal condition; Grade 1: horizontal growth; Grade 2: growth in vertical direction; Grade 3: significant growth in vertical direction; Grade 4: bridging with adjacent vertebrae. Redrawn from Rogers (1966) and Wada (1975).




Figure 3. Progressive stages of degenerative change of apophyseal joint surface. Grade 0: normal condition; Grade 1: osteophyte growing on the rim of articular surface without pitting; Grade 2: osteophyte growing on the rim of articular surface with lipping and slight pitting; Grade 3: osteophyte growing all around the rim of articular surface with moderate pitting; Grade 4: osteophyte significantly growing with severe pitting and unclear rim. Redrawn from Higuchi (1983).

according to the diagnostic criteria of Higuchi (1983). The grades were independently recorded for right and left sides of the superior and inferior articular processes of the respective vertebrae.

As a preliminary step toward analysis, the first author scored degenerative changes of Kumejima samples on independent occasions, and compared the data with the data scored by the third author to confirm that intra- and inter-observer errors were insignificant. We calculated crude prevalence rates of degenerative changes of the spine not per individual but per vertebra for comparisons of age-related changes, and sex-based and population-based differences because of the fragmentary condition of samples. We judged severity per vertebra using the most progressive stage of degenerative changes in the vertebral body and apophyseal joint. After that, we diagnosed greater than Grade 1 as a degenerative change and greater than Grade 3 as a severe degenerative change of the spine.

Fisher's exact probability test, χ^2 test and residual analysis were used to evaluate gender difference, age-related changes and regional differences. In addition, odds ratios (ORs) were calculated to evaluate age-related changes.

Results

Frequencies of degenerative changes per vertebra

Appendix Tables 1–4 show detailed basic data of observed and affected numbers, and frequencies of osteophytes (more than Grade 1) on the respective vertebral rims of vertebral bodies from the Okhotsk culture remains and me-

dieval Kamakura samples. Appendix Tables 5 and 6 also give basic data of observed and affected numbers and frequencies of degenerative changes of apophyseal joint (more than Grade 1) on the right and left sides of superior and inferior articular processes from the Okhotsk and Kamakura samples, respectively. Detailed data on the early-modern Kumejima samples have already been published (Moromizato et al., 2007).

Osteophytes on the vertebral body of the thoracic and lumbar vertebrae of each sex generally grew more anteriorly than posteriorly in both Okhotsk and Kamakura samples (Figure 4, Figure 5), as well as in the Kumejima samples (Moromizato et al., 2007); interestingly, osteophytes on the inferior rims on the posterior side of the upper cervical vertebrae (C2–C4) were more frequent than those on the anterior side in both sexes of the Kamakura series (Figure 5). Osteophytes on the vertebral body were more frequent in the lower cervical spine and the lower lumbar spine. Degenerative changes of apophyseal joint tended to be more pronounced in men of both series (Figure 6, Figure 7), and were more frequent in the lumbar vertebrae, especially at the levels of Th12–L1 and L4–L5, for the Okhotsk series. These frequencies corresponded to previous clinical findings (Resnick, 2002).

Age-related changes and sex-based differences in the Okhotsk and Kamakura series

In order to clarify the pattern of degenerative changes of joints, vertebrae were classified into four groups of cervical (C2–C7), upper thoracic (T1–T6), lower thoracic (T7–T12),





Figure 4. Frequencies of osteophytes on anterior and posterior sides of superior and inferior vertebral body surfaces: Okhotsk culture people. *Significantly different at the 0.05 level.

Figure 5. Frequencies of osteophytes on anterior and posterior sides of superior and inferior vertebral body surfaces: medieval Kamakura people. **, *Significantly different at the 0.01 or 0.05 level, respectively.

and lumbar vertebrae (L1-L5, S1) to calculate crude prevalence rates of osteophytes on the vertebral body and degenerative changes of apophyseal joint per vertebra. Age-related changes and sex-based differences in degenerative changes of the spine were then analyzed using the Okhotsk and Kamakura samples.

When the samples were divided into younger (under 40 years) and older groups (over 40 years), the older group of each sex had higher frequencies of osteophytes on the vertebral body and degenerative changes of apophyseal joint (more than Grade 1) than the younger group in the Kamakura sample ($P < 0.01$ or $P < 0.05$) except for degenerative changes of apophyseal joint of male cervical vertebrae, as shown in Table 2. Meanwhile, age-related differences of degenerative changes of apophyseal joint were not significant in males on the basis of the odds ratios. Severe degenerative changes (Grade 3 + Grade 4) were also more prevalent in the older group from Kamakura for each sex ($P < 0.01$ or $P < 0.05$) (Table 2).

In the Okhotsk series, on the other hand, age-related changes were not so clear, although more than half of the comparisons showed significant differences. For example, the differences in frequency between younger and older males were not statistically significant in all cases or in severe cases of degenerative changes of the apophyseal joint of lower thoracic vertebrae, and there were no significant differences between frequencies of all cases or in severe cases of degenerative changes of apophyseal joint of the upper thoracic region even in females (Table 3). In addition, severe osteophytes on the vertebral body of male cervical

vertebrae were more pronounced in the younger group in the Okhotsk samples ($P < 0.01$). Thus, severe degenerative changes of the spine appeared even in the younger generation of the Okhotsk people.

As for sex-based differences, the older female group in the Kamakura samples had a higher frequency of osteophytes on the body of the cervical vertebra than the older males ($P < 0.05$), whereas the older females had a lower frequency in the lower thoracic region ($P < 0.05$). The frequencies in younger males were significantly higher in the cervical, upper, and lower thoracic vertebrae than those in younger females ($P < 0.01$ or $P < 0.05$). Severe osteophytes on the vertebral body showed a few differences between sexes. Degenerative changes of apophyseal joint of all vertebral groups were more pronounced in younger males from Kamakura than in younger females ($P < 0.01$), while the frequency for upper thoracic vertebrae was only significantly higher in older males than in older females ($P < 0.05$). Although severe degenerative changes of apophyseal joint of the younger generation also tended to be affected in the same pattern, the frequency for lumbar vertebrae was higher in older females than in older males ($P < 0.01$).

In the Okhotsk series, younger males had a higher frequency of osteophytes on the body of cervical, upper, and lower thoracic spine than younger females ($P < 0.01$ or $P < 0.05$). However, as for the older generation, the frequency of osteophytes on the body of the cervical vertebrae was significantly higher in females than in males ($P < 0.05$), as was the case for the Kamakura series, while osteophytes on the body of upper thoracic vertebrae were more pronounced




Figure 6. Frequencies of degenerative changes on right and left sides of superior and inferior apophyseal joints: Okhotsk culture people. **, *Significantly different between sexes at the 0.01 or 0.05 level, respectively.




Figure 7. Frequencies of degenerative changes on right and left sides of superior and inferior apophyseal joints: medieval Kamakura people. **, *Significantly different between sexes at the 0.01 or 0.05 level, respectively.

in older males than in older females ($P < 0.01$). Severe osteophytes on the vertebral body in the older generation were more frequent in female cervical and male lumbar vertebrae ($P < 0.01$). Degenerative changes of apophyseal joint of cervical and lower thoracic spines were more pronounced in younger males than in younger females ($P < 0.01$), while the older generations exhibited no sex differences in this regard. Severe degenerative changes of apophyseal joint of lower thoracic vertebrae were more pronounced in younger males than in younger females ($P < 0.01$).

As a result, degenerative changes of the spine were generally more pronounced in males of both skeletal series, as described by Resnick (2002), while the degenerative changes of the spine of certain regions were more frequently seen in females. Because the younger males of both Kamakura and Okhotsk series tended to have high frequencies of degenerative changes of the spine, especially of apophyseal joint, age-related changes in males were obscure on the basis of the results of Fisher's exact probability tests and odds ratios.

Population-based differences

Next, we evaluated population differences of degenerative changes of the spine among the Okhotsk, medieval Kamakura, and early-modern Kumejima series (Table 4, Table 5). Although the three series were least affected in the upper thoracic vertebrae, with the exception of Okhotsk males ($P < 0.01$ or $P < 0.05$), they clearly showed different patterns of osteophytes on the vertebral body. For example,

a significantly high frequency of affected cervical vertebrae was found for the Okhotsk people ($P < 0.01$ or $P < 0.05$), while the Kumejima samples had the highest frequency in the lumbar vertebrae ($P < 0.01$). As mentioned previously (Moromizato et al., 2007), the Kumejima females had a higher frequency in the cervical vertebrae than males ($P < 0.01$). In the Kamakura series, males were most affected in the lower thoracic vertebrae ($P < 0.01$). Among the three series, the Kumejima samples had the highest frequency in the lumbar vertebrae ($P < 0.01$).

As for severe osteophytes on the vertebral body, all three series had the highest frequencies in the lumbar vertebrae, with the exception of the Okhotsk females ($P < 0.01$). Severe osteophytes on the body of other vertebral parts displayed markedly different patterns. The Okhotsk series had higher frequencies on the cervical vertebrae than the other two series ($P < 0.01$ or $P < 0.05$), while the Kamakura series generally had higher frequencies of severe osteophytes on the vertebral body, which was contrary to our expectations. The Kumejima series was the least affected in the cervical and thoracic vertebrae among the three series ($P < 0.01$).

Affected patterns of degenerative changes of apophyseal joint within populations showed clearer differences. The Okhotsk series of each sex had the lowest frequencies in the cervical vertebrae and the highest frequencies in the lumbar vertebrae ($P < 0.01$ or $P < 0.05$), which contrasted with the highest frequency in the cervical vertebrae in the Kumejima series ($P < 0.01$). The Kamakura series of each sex had

Table 2. Comparisons of crude prevalence rates of degenerative changes of spines in medieval Kamakura

	Female			Male		
	A/O	%		A/O	%	
Osteophytes on the vertebral body						
Cervical vertebrae	Older	105/154	68.2	>	67/122	54.9
	Younger	51/199	25.6	<<	90/221	40.7
	OR (95% CI)	2.01 (1.20–3.39)	**		1.77 (1.13–2.77)	**
Upper thoracic vertebrae	Older	64/127	50.3		57/102	55.9
	Younger	47/201	23.4	<	76/243	31.3
	OR (95% CI)	3.33 (2.07–5.36)	**		2.78 (1.73–4.48)	**
Lower thoracic vertebrae	Older	83/137	60.6	<	85/118	72.0
	Younger	63/202	31.2	<	110/257	42.8
	OR (95% CI)	3.39 (2.15–5.34)	**		3.44 (2.15–5.52)	**
Lumbar vertebrae (+S1)	Older	112/142	78.9		89/123	72.4
	Younger	56/233	24.0		69/257	26.8
	OR (95% CI)	11.8 (7.14–19.5)	**		7.13 (4.40–11.55)	**
Severe osteophytes on the vertebral body (Grade 3 + Grade 4)						
	A/O	%		A/O	%	
Cervical vertebrae	Older	28/154	18.2		21/122	17.2
	Younger	1/199	0.5	<	8/221	3.6
	OR (95% CI)	44.0 (5.91–327.4)	**		5.54 (2.37–12.9)	**
Upper thoracic vertebrae	Older	13/127	10.2	>>	1/102	1.0
	Younger	2/201	1.0		9/243	3.7
	OR (95% CI)	11.3 (2.52–51.2)	**		0.26 (0.03–2.06)	
Lower thoracic vertebrae	Older	23/137	16.8		17/118	14.4
	Younger	9/202	4.5		11/257	4.3
	OR (95% CI)	4.33 (1.94–9.67)	**		3.76 (1.70–8.32)	**
Lumbar vertebrae (+S1)	Older	68/142	47.8		44/123	35.8
	Younger	12/233	5.1		13/257	5.1
	OR (95% CI)	16.9 (8.68–33.0)	**		10.5 (5.36–20.4)	**
Degenerative changes of apophyseal joint						
	A/O	%		A/O	%	
Cervical vertebrae	Older	39/168	23.2		20/123	16.3
	Younger	3/228	1.3	<<	30/246	12.2
	OR (95% CI)	22.4 (6.78–73.8)	**		1.39 (0.76–2.58)	
Upper thoracic vertebrae	Older	22/113	19.5	<	28/86	32.6
	Younger	10/192	5.2	<<	49/235	20.9
	OR (95% CI)	4.40 (2.00–9.68)	**		1.64 (0.94–2.84)	*
Lower thoracic vertebrae	Older	43/126	34.1		41/101	40.6
	Younger	15/190	7.9	<<	63/242	26.0
	OR (95% CI)	6.04 (3.18–11.5)	**		1.94 (1.19–3.17)	**
Lumbar vertebrae (+S1)	Older	51/126	40.4		33/107	30.8
	Younger	24/219	11.0	<<	55/251	21.9
	OR (95% CI)	5.53 (3.18–9.61)	**		1.59 (0.96–2.64)	*

Table 2. (continued)

	Female		Male		
	A/O	%	A/O	%	
Severe degenerative changes of apophyseal joint (Grade 3 + Grade 4)					
Cervical vertebrae	Older	18/168	10.7	11/123	8.9
	Younger	1/228	0.4	2/246	0.8
	OR (95% CI)	27.2 (3.60–206.2)	**	12.0 (2.61–55.0)	**
Upper thoracic vertebrae	Older	6/113	5.3	11/86	12.8
	Younger	1/192	0.5	9/235	3.8
	OR (95% CI)	10.7 (1.27–90.1)	*	3.68 (1.47–9.23)	**
Lower thoracic vertebrae	Older	15/126	11.9	10/101	9.9
	Younger	0/190	0.0	11/242	4.5
	OR (95% CI)	–	**	2.31 (0.95–5.62)	*
Lumbar vertebrae (+S1)	Older	26/126	20.6	8/107	7.5
	Younger	3/219	1.4	13/251	5.2
	OR (95% CI)	18.7 (5.54–63.3)	**	1.48 (0.59–3.68)	

** , *Significantly different between younger and older at the 0.01 or 0.05 level.

<<, >>, <, > Significantly different between male and female at the 0.01 or 0.05 level.

A, number of vertebrae showing lesions (affected); O, number of vertebrae actually observed; F: frequency (%).

OR: odds ratio; 95% CI: 95% confidence interval.

generally low frequencies of degenerative changes of apophyseal joint, especially in the cervical and upper thoracic regions, because of the relatively young age distribution.

Among the three series, severe degenerative changes of apophyseal joint dominantly affected the Okhotsk series of each sex. The Kumejima samples showed low frequencies ($P < 0.01$ or $P < 0.05$) although the relatively high frequencies in the cervical vertebrae remained ($P < 0.01$).

Discussion

We confirmed that degenerative changes of the spine in ancient populations were more pronounced in men and older groups, as in contemporary populations. In addition, we revealed clear population-based and sex-based differences among the ancient groups from the Japanese archipelago.

Osteophytes on the vertebral body

Osteophytes developed more on the anterior side of the vertebral body in the three ancient peoples and recent Japanese (Wada, 1975). Age-related changes were clearly seen in the medieval Kamakura, early-modern Kumejima (Moromizato et al., 2007), and recent Japanese (Wada, 1975).

Osteophytes on the vertebral body tend to be more pronounced in men, leading to clinical cases (Resnick, 2002). However, older females had a higher frequency in the cervical vertebrae than the older males of both Kamakura and Okhotsk series. In addition, the Kumejima females of both younger and older groups combined had a higher frequency in the cervical vertebrae than males (Moromizato et al., 2007). Because there were similar frequencies between women and men and degenerative changes are more severe in men in clinical medicine (Resnick, 2002), this gender dif-

ference may be caused by some dynamic factors, including abnormal forces on the spinal column (Shedid and Benzel, 2007). Moromizato et al. (2007) mentioned that the Ryukyu women carry heavy loads on their head, called 'Kasami.' Although we do not know about the existence of such customs in the medieval Kamakura and Okhotsk periods, findings in other archeological cases suggest that older females followed this practice (Lovell, 1994).

Osteophytes on the body of the lumbar vertebrae were most frequent in the Kumejima samples, while those of the lower thoracic vertebrae and the cervical vertebrae were most frequent in the medieval Kamakura and Okhotsk series, respectively. Degenerative changes of the lumbar spine were found to be frequent in contemporary and archeological populations, including the Canadian Inuit (Merbs, 1983; Bridges, 1994; Larsen, 1997; Resnick, 2002; and many others). In Japan, the lumbar spine was found to be most affected in the Jomon, Yayoi, early-modern Edo and recent Japanese samples (Wada, 1975; Suzuki, 1978; Fukushima, 1988). Severe osteophytes on the body of the lumbar vertebrae were most frequent not only in the Kumejima series but also in the medieval Kamakura samples. Thus, the Okhotsk series had the peculiar characteristic of a higher frequency of osteophytes on the body of the cervical spine among populations from the Japanese archipelago, although the Kumejima females had the highest frequency, as mentioned above. The cervical vertebrae only support loads of or on the head, while the head, trunk and upper extremities produce loads on the lumbar vertebrae (Roh et al., 2005). Although we know of no relevant customs in prehistoric Okhotsk culture, some dynamic loads caused a high frequency of degenerative changes of cervical spine of clinical severity in other archeological cases (Shedid and Benzel, 2007; Lovell, 1994).

Table 3. Comparisons of crude prevalence rates of degenerative changes of spines in the Okhotsk series

		Female		Male		
		A/O	F	A/O	F	
Osteophytes on the vertebral body						
Cervical vertebrae	Older	46/75	61.3	>	19/47	40.4
	Younger	12/49	24.5	<<	37/64	57.8
	OR (95% CI)	4.89 (2.20–10.9)	**		0.50 (0.23–1.06)	
Upper thoracic vertebrae	Older	20/80	25.0	<<	26/51	51.0
	Younger	1/39	2.6	<<	19/73	26.0
	OR (95% CI)	12.7 (1.63–98.3)	**		2.96 (1.38–6.31)	**
Lower thoracic vertebrae	Older	42/83	50.6		35/53	66.0
	Younger	1/43	2.3	<	11/86	12.8
	OR (95% CI)	43.0 (5.65–327.4)	**		13.3 (5.66–13.0)	**
Lumbar vertebrae (+S1)	Older	31/68	45.6	<	33/50	66.0
	Younger	8/48	16.7		16/81	19.8
	OR (95% CI)	4.19 (1.71–10.3)	**		7.89 (3.54–17.6)	**
Severe osteophytes on the vertebral body (Grade 3 + Grade 4)						
		A/O	F		A/O	F
Cervical vertebrae	Older	15/75	20.0	>>	1/47	2.1
	Younger	2/49	4.1	<	12/64	18.8
	OR (95% CI)	5.88 (1.28–27.0)	**		0.09 (0.01–0.75)	**
Upper thoracic vertebrae	Older	3/80	3.8		5/51	9.8
	Younger	0/49	0.0		2/73	2.7
	OR (95% CI)	—			3.86 (0.72–20.7)	*
Lower thoracic vertebrae	Older	7/83	8.4		4/53	7.5
	Younger	0/43	0.0		6/86	7.0
	OR (95% CI)	—	**		1.09 (0.29–4.05)	
Lumbar vertebrae (+S1)	Older	10/68	14.7	<<	20/50	40.0
	Younger	3/48	6.3		8/80	10.0
	OR (95% CI)	2.59 (0.67–9.95)	*		6.00 (2.38–15.1)	**
Degenerative changes of apophyseal joint						
		A/O	F		A/O	F
Cervical vertebrae	Older	26/87	29.9		16/41	39.0
	Younger	4/61	6.6	<<	20/82	24.4
	OR (95% CI)	6.07 (2.00–18.5)	**		1.98 (0.89–4.44)	
Upper thoracic vertebrae	Older	30/75	40.0		29/56	51.8
	Younger	19/52	36.5		20/80	25.0
	OR (95% CI)	1.16 (0.56–2.40)			3.22 (1.56–6.68)	**
Lower thoracic vertebrae	Older	35/76	46.1		23/53	43.4
	Younger	7/53	13.2	<<	33/87	37.9
	OR (95% CI)	5.61 (2.25–14.0)	**		1.25 (0.63–2.51)	
Lumbar vertebrae (+S1)	Older	45/81	55.6		33/55	60.0
	Younger	14/54	25.9		28/79	35.4
	OR (95% CI)	3.57 (1.69–7.56)	**		2.73 (1.34–5.56)	**

Table 3. (continued)

	Female		Male		
	A/O	F	A/O	F	
Severe degenerative changes of apophyseal joint (Grade 3 + Grade 4)					
Cervical vertebrae	Older	13/87	14.9	9/57	15.8
	Younger	2/61	3.3	5/82	6.1
	OR (95% CI)	5.18 (1.13–23.9)	*	2.87 (0.91–9.13)	
Upper thoracic vertebrae	Older	9/75	12.0	10/56	17.9
	Younger	4/52	7.7	4/80	5.0
	OR (95% CI)	1.64 (0.48–5.63)		4.13 (1.22–13.9)	*
Lower thoracic vertebrae	Older	20/76	26.3	11/53	20.8
	Younger	2/53	3.8	20/87	23.0
	OR (95% CI)	9.11 (2.03–40.9)	**	0.88 (0.38–2.01)	
Lumbar vertebrae (+S1)	Older	21/81	25.9	11/55	20.0
	Younger	4/54	7.4	9/79	11.4
	OR (95% CI)	4.38 (1.41–13.6)	*	1.94 (0.75–5.07)	*

** , * : Significantly different between younger and older at the 0.01 or 0.05 level.

<<, >>, <, > Significantly different between male and female at the 0.01 or 0.05 level.

A, number of vertebrae showing lesions (affected); O, number of vertebrae actually observed; F: frequency (%).

OR: odds ratio; 95% CI: 95% confidence interval.

Degenerative changes of apophyseal joint

As mentioned above, the bodies of vertebrae are connected by intervertebral discs (symphyses), while the apophyseal joints are paired, true synovial joints. Therefore, the degenerative processes of the two joints may occur as independent phenomena (Resnick, 2002). However, intervertebral disc degeneration (intervertebral chondrosis and spondylosis deformans) causes narrowing of the intervertebral disc space, thus probably resulting in apophyseal articular alterations (Kalichman and Hunter, 2007). Although we could not analyze direct correlations between osteophytes on the vertebral body and degenerative changes of apophyseal joint within individuals because of the fragmented nature of skeletal materials, the frequency patterns of the samples were compared.

The lower lumbar vertebrae were most affected in the Okhotsk series, coinciding with findings for contemporary people (Fujiwara et al., 1999), whereas the Kamakura and Kumejima samples showed no clear peak in the lumbar vertebrae. Age-related differences in degenerative changes of apophyseal joint were more obscure than those in osteophytes on the vertebral body, especially in the Okhotsk males, whereas the recent Japanese showed clear age-related changes (Higuchi, 1983). This may be due to other factors, including labor and other activity-related causes.

Degenerative changes of the apophyseal joint also tend to be more pronounced in men, as in clinical cases (Resnick, 2002). It is interesting that the frequency of severe degenerative changes of apophyseal joint in the lumbar vertebrae of the Kamakura series was higher in older females than in older males. Because postmenopausal women had a high frequency of degenerative changes of lumbar apophyseal joint owing to increased expression of estrogen receptor (Ha et

al., 2005), this high frequency seen in the Kamakura older females was possibly caused by age-related changes.

We found that there were marked differences in the pattern of effects exhibited between the Okhotsk and Kumejima series, i.e. the highest frequency for the lumbar spine in the Okhotsk series contrasts well with that of the cervical spine in the Kumejima series. However, an inverse pattern was exhibited for the frequencies of osteophytes on the vertebral body.

This may be explained by differences of anatomy and biomechanics between the apophyseal joints of cervical and lumbar vertebrae. The line of gravity runs just anterior to the apophyseal joints in the cervical vertebrae, while the line runs within the bodies of lumbar vertebrae, separate from the apophyseal joints (Standring, 2008). The superior and inferior flat articular facets of the cervical spine face posterosuperiorly and anteroinferiorly, respectively, at an angle of about 45° to the transverse plane at the upper cervical spine (Aiello and Dean, 1990; Kirpalani and Mitra, 2008). On the other hand, the facet joints of the lumbar vertebrae are very distinctive (Aiello and Dean, 1990; Kalichman and Hunter, 2007). The concave superior and convex inferior articular facets face medially and laterally, respectively, making an interlocking joint. Thus, the cervical apophyseal joint bears more weight and rotates easily, while the lumbar apophyseal joint has restricted rotation and prohibits anterior and lateral sliding.

The higher frequencies of degenerative changes of cervical apophyseal joint in the Kumejima series could be explained by additional loads on the head, especially in females (Moromizato et al., 2007). However, we would like to infer several dynamic reasons for the higher frequency of degenerative changes of lumbar apophyseal joint in the

Table 4. Comparisons of crude prevalence rates of osteophytes on the vertebral body in the Okhotsk, Kamakura and Kumejima series

Osteophytes on the vertebral body		Female		Male	
		A/O	F	A/O	F
Cervical vertebrae	Okhotsk	58/124	46.8##	56/111	50.5#
	Kamakura	156/353	44.2**	157/343	45.8
	Kumejima	138/238	58.0**	143/316	45.3##
		**		>>	
Upper thoracic vertebrae	Okhotsk	21/119	17.6***	45/124	36.3
	Kamakura	111/328	33.8##	133/345	38.6#
	Kumejima	79/208	38.0***	113/306	36.9##
		**		<<	
Lower thoracic vertebrae	Okhotsk	43/126	34.1**	46/139	33.1**
	Kamakura	152/339	44.8	195/375	52.0##
	Kumejima	114/208	54.8**	191/298	64.1***
		**		<	
Lumbar vertebrae (+S1)	Okhotsk	39/116	33.6**	49/131	37.4**
	Kamakura	168/375	44.8**	158/380	42.1**
	Kumejima	172/241	71.4***	236/318	74.2***
		**		**	
Severe osteophytes on the vertebral body (Grade 3 + Grade 4)					
		A/O	F	A/O	F
Cervical vertebrae	Okhotsk	17/124	13.7***	13/111	11.7*
	Kamakura	29/353	8.2#	29/343	8.5*
	Kumejima	7/238	2.9***	8/316	2.5***
		**		**	
Upper thoracic vertebrae	Okhotsk	3/119	2.5##	7/124	5.6***
	Kamakura	15/328	4.6***	10/345	2.9##
	Kumejima	0/208	0.0***	0/306	0.0***
		**		**	
Lower thoracic vertebrae	Okhotsk	7/126	5.6	10/139	7.2
	Kamakura	32/339	9.4**	28/375	7.5*
	Kumejima	3/208	1.4***	8/298	2.7***
		**		*	
Lumbar vertebrae (+S1)	Okhotsk	13/116	11.2	28/131	21.4##
	Kamakura	80/375	21.3##	57/380	15.0##
	Kumejima	49/241	20.3##	60/318	18.9##
		<		>	

** , * : Significantly different among three samples at the 0.01 or 0.05 level.

##, # : Significantly different among vertebral regions within the respective skeletal series at the 0.01 or 0.05 level.

<<, >>, <, > : Significantly different between male and female at the 0.01 or 0.05 level.

A, number of vertebrae showing lesions (affected); O, number of vertebrae actually observed; F: frequency (%).

Okhotsk series because severe patterns were exclusively present at high frequency in this series among the three samples.

People from Okhotsk culture

The Okhotsk culture appeared in southern Sakhalin and spread to northeastern Hokkaido and the Kuril Islands from the 5th to 12th century AD (Yamaura and Ushiro, 1999; Amano, 2003). The average annual temperature today is about 7°C, with -5°C in January, at Wakkanai City, the

northernmost city on Hokkaido. Sea drift ice occupies the coast of the Okhotsk Sea from January to March. The Okhotsk culture developed considerable sea-mammal hunting, in both shallow and deep water, unlike the native population of Hokkaido (Hudson, 2004). These sea mammals consisted of fur seal, whales, sea lions and others. For hunting and fishing, the Okhotsk people used bone tools, including hooks and harpoons, as well as iron tools (Amano, 2003). In addition, they traveled in boats and probably fished using nets based on findings of stone weights. The

Table 5. Comparisons of crude prevalence rates of degenerative changes of apophyseal joint in the Okhotsk, Kamakura and Kumejima series.

Degenerative changes of apophyseal joint		Female		Male	
		A/O	F	A/O	F
Cervical vertebrae	Okhotsk	30/148	20.3##	36/139	25.9##
	Kamakura	42/396	10.6***	45/378	11.9***
	Kumejima	132/276	47.8***	169/361	46.8***
		**		**	
Upper thoracic vertebrae	Okhotsk	49/127	38.6**	49/136	36.0**
	Kamakura	32/305	10.5***	<< 66/323	20.4*
	Kumejima	25/200	12.5##	<< 68/300	22.7##
		**		**	
Lower thoracic vertebrae	Okhotsk	42/129	32.6*	56/140	40.0
	Kamakura	58/316	18.4**	<< 97/354	27.4***
	Kumejima	62/207	30.0**	107/295	36.3
		**		**	
Lumbar vertebrae (+S1)	Okhotsk	59/135	43.7***	61/134	45.5**
	Kamakura	75/345	21.7***	81/373	21.7**
	Kumejima	91/236	38.6**	126/312	40.4**
		**		**	
Severe degenerative changes of apophyseal joint (Grade 3 + Grade 4)					
		A/O	F	A/O	F
Cervical vertebrae	Okhotsk	15/148	10.1*	14/139	10.1*
	Kamakura	19/396	4.8	13/378	3.4*
	Kumejima	14/276	5.1##	23/361	6.4##
		*		*	
Upper thoracic vertebrae	Okhotsk	13/127	10.2**	14/136	10.3**
	Kamakura	7/305	2.3#	< 19/323	5.9
	Kumejima	0/200	0.0***	1/300	0.3***
		**		**	
Lower thoracic vertebrae	Okhotsk	22/129	17.1**	31/140	22.1***
	Kamakura	15/316	4.7	18/354	5.1
	Kumejima	2/207	1.0**	5/295	1.7**
		**		**	
Lumbar vertebrae (+S1)	Okhotsk	25/135	18.5**	20/134	14.9**
	Kamakura	29/345	8.4##	> 14/373	3.8
	Kumejima	5/236	2.1**	8/312	2.6**
		**		**	

**, *: Significantly different among three samples at the 0.01 or 0.05 level.

##, #: Significantly different among vertebral regions within the respective skeletal series at the 0.01 or 0.05 level.

<<, >>, <, > Significantly different between male and female at the 0.01 or 0.05 level.

A, number of vertebrae showing lesions (affected); O, number of vertebrae actually observed; F: frequency (%).

nature of subsistence of the Okhotsk culture was recently confirmed using stable isotope analysis (Naito et al., 2010).

There has been little bioarcheological research on the Okhotsk people; Kodama (1948) mentioned severe dental wear and Yamaguchi (1995) identified a wound caused by a stone arrowhead in the right hip bone of a male specimen. Recently, oral health has attracted attention (Ishida et al., 1994; Hudson, 2004; Fukumoto et al., 2007; Oxenham and Matsumura, 2008; Hoover and Matsumura, 2008). However, only vertebral compression fractures were reported

with regard to the skeletal system of the Okhotsk people (Ishida et al., 1994; Ishida and Matsumura, 2000).

Degenerative changes were found even in the younger Okhotsk individuals. As mentioned above, degenerative changes of lumbar apophyseal joint were more pronounced in the Okhotsk people. This was possibly caused by travel by boat and net fishing, as suggested by archeological evidence, because of the associated need for much rotation, extension and flexion of the lumbar spine.

Medieval Kamakura and early-modern Kumejima peoples

Intensive paleodemographic research has been performed and has revealed that the medieval Kamakura people were not long lived (Nagaoka et al., 2006; Nagaoka and Hirata, 2008). In addition, many weapon-related traumas were found among the skeletal remains from medieval Kamakura (Suzuki et al., 1956; Hirata et al., 2004; Nagaoka et al., 2009, 2010). However, there have been few studies on the life activity patterns of these people. As mentioned above, the skeletal remains from medieval Kamakura were thought to be from common people (Hirata et al., 2004; Nagaoka et al., 2006). Residents in the medieval city of Kamakura had many different occupations, such as merchants, civil servants, workmen and housekeepers. Although they had no specific labor patterns, age-related changes of degenerative diseases were apparent, as in contemporary and clinical cases (Wada, 1975).

Regarding the Ryukyu Islands, there have been only a few bioarcheological studies. Dental disease, including dental caries and linear enamel hypoplasia, was investigated by Oyamada et al. (1996) and Hudson and Takamiya (2003). Degenerative changes in the elbow joints and lumbar vertebrae were found in early-modern samples on Ishigaki Island, in the southernmost island group of the Ryukyu Islands (Zukeran et al., 2002). As for the early-modern Kumejima samples, dental diseases were reported to have been found with a high rate of dental caries in adult females, suggesting differences in food preference may have led to this sex difference, as suggested by isotopic analysis (Irei et al., 2008). We have previously reported that the pattern of a high frequency of degenerative changes in the lumbar spine coincided with that of farming peoples, including the prehistoric northern Thailand people from the Ban Chiang site (Pietruszewsky and Douglas, 2002), which is due to lasting flexion posture during farming, and that the gender difference seen in the cervical spine may have been caused by labor specific to women, such as carrying items on the head (Moromizato et al., 2007).

Degenerative joint diseases have been investigated in clinical medicine (Mukai et al., 2009a, b). These established epidemiological studies using very large populations and exact diagnosis revealed many risk factors and the etiology of degenerative joint diseases. Because archeological human skeletal remains are imperfect for age and sex determination, in terms of preservation, population size, and especially lack of soft tissue, methods and results of analysis cannot guarantee definitive conclusions.

For example, there are some limitations to this study. First, because the age distribution in the Okhotsk females tended to be biased towards an older age than in the males ($P = 0.15$), male dominant sex differences almost disappeared when conjoined with age groups. Second, the age distribution of the Kumejima people was different from those of the other peoples owing to their relatively long life expectancy. Therefore, there was a good possibility that the high frequencies of degenerative changes seen in the Kumejima series were caused not by population-specific events but were merely aging effects.

However, in this study using relatively large populations

of three ancient human groups, we could reveal age-related, sex-based, and population-based patterns of degenerative joint changes among peoples of the Japanese archipelago. We will continue to investigate such skeletal changes of many other ancient peoples in order to obtain important new information on their quality of life.

Acknowledgments

This study was supported in part by Grants-in-aid for Scientific Research from the Japan Society for the Promotion of Science (Nos. 18370099, 22370087). We are deeply grateful to two anonymous reviewers for their valuable comments.

References

- Aiello L. and Dean C. (1990) *An Introduction to Human Evolutionary Anatomy*. Academic Press, London.
- Amano T. (2003) What is the Okhotsk culture? In: Nomura T. and Utagawa Y. (eds.), *Epi-jomon, Okhotsk Culture*. Hokkaido Shinbunsha, Sapporo, pp. 110–133 (in Japanese).
- Bridges P.S. (1994) Vertebral arthritis and physical activities in the prehistoric southeastern United States. *American Journal of Physical Anthropology*, 93: 83–93.
- Coggon D., Kellingray S., Inskip H., Croft P., Campbell L., and Cooper C. (1998) Osteoarthritis of the hip and occupational lifting. *American Journal of Epidemiology*, 147: 523–528.
- Dieppe P.A. and Lohmander L.S. (2005) Pathogenesis and management of pain in osteoarthritis. *Lancet*, 365: 965–973.
- Felson D.T., Lawrence R.C., Dieppe P.A., Hirsh R., Helmick C.G., Jordan J.M., Kington R.S., Lane N.E., Nevitt M.C., Zhang Y., Sowers M., McAlindon T., Spector T.D., Poole A.R., Yanovski S.Z., Ateshian G., Sharma L., Buckwalter J.A., Brandt K.D., and Fries J.F. (2000) Osteoarthritis: new insights. Part 1: The disease and its risk factors. *Annals of Internal Medicine*, 133: 635–646.
- Freund M. and Sartor K. (2006) Degenerative spine disorders in the context of clinical findings. *European Journal of Radiology*, 58: 15–26.
- Fujiwara A., Tamai K., Yamato M., An H.S., Yoshida H., Saotome K., and Kurihashi A. (1999) The relationship between facet joint osteoarthritis and disc degeneration of the lumbar spine: an MRI study. *European Spine Journal*, 8: 396–401.
- Fukumine T., Doi N., Ishida H., Zukeran C., Sensui S., Saso A., and Higa T. (2001) Human skeletal remains from the Yacchino-Gama and Kanjinbaru grave sites. In: Okinawa Prefectural Archaeological Center (ed.), *Yacchi-no-Gama, Kanjinbaru Grave Sites*. Okinawa Prefectural Archaeological Center Research Report No. 6, Nishihara, pp. 345–385 (in Japanese).
- Fukumine T., Hanihara T., Nishime A., and Ishida H. (2006) Non-metric cranial variation of early modern human skeletal remains from Kumejima, Okinawa and the peopling of the Ryukyu Islands. *Anthropological Science*, 114: 141–151.
- Fukumoto I., Amano T., Ono H., Matsumura H., Yoneda M., and Ishida H. (2007) Prevalences of dental enamel hypoplasia and cribra orbitalia in the human crania associated with the Okhotsk culture. *Anthropological Science*, 115: 255.
- Fukushima K. (1988) On the lesion of bones of Yayoi people in Southwest Japan. *Fukuoka Acta Medica*, 79: 227–248.
- Gallucci M., Puglielli E., Splendiani A., Pistoia F., and Spacca G. (2005) Degenerative disorders of the spine. *European Radiology*, 15: 591–598.
- Ha K., Chang C., Kim K., Kim Y., Na K., and Lee J. (2005) Expression of estrogen receptor of the facet joints in degenerative spondylolithesis. *Spine*, 30: 562–566.
- Higuchi Y. (1983) A study on the morphological changes in the human cervical apophyseal joints with increasing age on

- macerated skeletons. *Sapporo Medical Journal*, 52: 181–204.
- Hirata K., Nagaoka T., and Hoshino K. (2004) Analysis of injuries by swords in medieval Japanese skeletons from Yuigahama, Kamakura. *Anthropological Science (Japanese series)*, 112: 19–26.
- Hoover K.C. and Matsumura H. (2008) Temporal variation and interaction between nutritional and developmental instability in prehistoric Japanese populations. *American Journal of Physical Anthropology*, 137: 469–478.
- Hudson M.J. (2004) The perverse realities of change: world system incorporation and the Okhotsk culture of Hokkaido. *Journal of Anthropological Archaeology*, 23: 290–308.
- Hudson M.J. and Takamiya H. (2003) Dental pathology and subsistence change in late prehistoric Okinawa. *Indo-Pacific Prehistory Association Bulletin*, 21: 68–76.
- Irei K., Doi N., Fukumine T., Nishime A., Hanihara T., Yoneda M., and Ishida H. (2008) Dental diseases of human skeletal remains from the early-modern period of Kumejima Island, Okinawa, Japan. *Anthropological Science*, 116: 149–159.
- Ishida H. (1996) Metric and nonmetric cranial variation of the prehistoric Okhotsk people. *Anthropological Science*, 104: 233–258.
- Ishida H. and Matsumura H. (2000) Human skeletal remains of the Okhotsk culture recovered at Hamanaka 2 site, Rebun-cho, Hokkaido in 1994–95. In: Nishimoto T. (ed.), Report on the Research Excavation at the Site of Hamanaka 2. *Bulletin of the National Museum of Japanese History*, 85: 264–270.
- Ishida H., Hanihara T., Kondo O., and Ohshima N. (1994) A human skeleton of the early phase of the Okhotsk culture unearthed at the Hamanaka-2 site, Rebun Island, Hokkaido. *Anthropological Science*, 102: 363–378.
- Jurmain R. (2000) Degenerative joint disease in African great apes: an evolutionary perspective. *Journal of Human Evolution*, 39: 185–203.
- Kalichman L. and Hunter D.J. (2007) Lumbar facet joint osteoarthritis: a review. *Seminars in Arthritis and Rheumatism*, 37: 69–80.
- Kirpalani D. and Mitra R. (2008) Cervical facet joint dysfunction: a review. *Archives of Physical Medicine and Rehabilitation*, 89: 770–774.
- Kodama S. (1948) Moyoro Shell Heap. Hokkaido Genshi Bunka Kenkyukai, Sapporo (in Japanese).
- Komesu A., Hanihara T., Amano T., Ono H., Yoneda M., Dodo Y., Fukumine T., and Ishida H. (2008) Nonmetric cranial variation in human skeletal remains associated with Okhotsk culture. *Anthropological Science*, 116: 33–47.
- Larsen C.S. (1997) *Bioarchaeology: Interpreting Behavior from the Human Skeleton*. Cambridge University Press, Cambridge.
- Lieverse A.R., Weber A.W., Bazaliiskiy V.I., Goriunova O.I., and Savel'ev N.A. (2007) Osteoarthritis in Siberia's Cis-Baikal: skeletal indicators of hunter-gatherer adaptation and cultural change. *American Journal of Physical Anthropology*, 132: 1–16.
- Lovell N.C. (1994) Spinal arthritis and physical stress at Bronze Age Harappa. *American Journal of Physical Anthropology*, 93: 149–164.
- Merbs C.F. (1983) Patterns of activity-induced pathology in a Canadian Inuit population. *Archaeological Survey of Canada Paper*, 119: 1–119.
- Moromizato K., Fukumine T., Doi N., Hanihara T., Nishime A., Yoneda M., and Ishida H. (2007) Degenerative diseases of the spines of early modern human remains from Kumejima, Okinawa. *Anthropological Science (Japanese series)*, 115: 25–36.
- Mukai S., Akune T., Oka H., Mabuchi A., En-yo Y., Yoshida M., Saika A., Nakamura K., Kawaguchi H., and Yoshimura N. (2009a) Association of occupational activity with radiographic knee osteoarthritis and lumbar spondylosis in elderly patients of population-based cohorts: a large-scale population-based study. *Arthritis and Rheumatism*, 61: 779–786.
- Mukai S., Oka H., Akune T., Mabuchi A., En-yo Y., Yoshida M., Saika A., Suzuki T., Yoshida H., Ishibashi H., Yamamoto S., Nakamura K., Kawaguchi H., and Yoshimura N. (2009b) Prevalence of radiographic lumbar spondylosis and its association with low back pain in elderly subjects of population-based cohorts: the ROAD study. *Annals of the Rheumatic Diseases*, 68: 1401–1406.
- Nagaoka T. and Hirata K. (2008) Demographic structure of skeletal populations in historic Japan: a new estimation of adult age-at-death distributions based on the auricular surface of the ilium. *Journal of Archaeological Science*, 35: 1370–1377.
- Nagaoka T., Hirata K., Yokota E., and Matsumura S. (2006) Paleodemography of a medieval population in Japan: analysis of human skeletal remains from the Yuigahama-minami site. *American Journal of Physical Anthropology*, 131: 1–14.
- Nagaoka T., Uzawa K., and Hirata K. (2009) Weapon-related traumas of human skeletons from Yuigahama Chusei Shudan Bochi, Japan. *Anatomical Science International*, 84: 170–181.
- Nagaoka T., Uzawa K., and Hirata K. (2010) Evidence of weapon-related traumas in medieval Japan: observations of the human crania from Seiyokan. *Anthropological Science*, 118: 129–140.
- Naito Y.I., Chikaraishi Y., Ohkouchi N., Mukai H., Shibata Y., Honch N.V., Dodo Y., Ishida H., Amano T., Ono H., and Yoneda M. (2010) Dietary reconstruction of the Okhotsk Culture of Hokkaido, Japan, based on nitrogen isotopic composition of amino acids: implication for the correction of radiocarbon marine reservoir effects on human bones. *Radiocarbon*, 52: 671–681.
- Oyamada J., Manabe Y., Kitagawa Y., and Rokutanda A. (1996) Dental morbid condition of hunter-gatherers on Okinawa island during the middle period of the prehistoric shell midden culture and of agriculturalists in northern Kyushu during the Yayoi period. *Anthropological Science*, 104: 261–280.
- Oxenham M.F. and Matsumura H. (2008) Oral and physiological paleohealth in cold adapted peoples: Northeast Asia, Hokkaido. *American Journal of Physical Anthropology*, 135: 64–74.
- Pietrusewsky M. and Douglas M.T. (2002) Ban Chiang: A Prehistoric Village Site in Northeast Thailand I: The Human Skeletal Remains. University of Pennsylvania Press, Philadelphia.
- Pytel P., Wollmann R.L., Fessler R.G., Krausz T.N., and Montag A.G. (2006) Degenerative spine disease. Pathologic findings in 985 surgical specimens. *American Journal of Clinical Pathology*, 125: 193–202.
- Resnick D. (2002) Degenerative disease of the spine. In Resnick D. (ed.), *Diagnosis of Bone and Joint Disorders*. W.B. Saunders Co., Philadelphia, pp. 1382–1475.
- Rogers S.L. (1966) The need for a better means of recording pathological bone proliferation in joint areas. *American Journal of Physical Anthropology*, 25: 171–176.
- Roh J.S., Teng A.L., Yoo J.U., Davis J., Furey C., and Bohlman H.H. (2005) Degenerative disorders of the lumbar and cervical spine. *Orthopedic Clinics of North America*, 36: 255–262.
- Rojas-Sepúlveda C., Ardagna Y., and Dutour O. (2008) Paleoepidemiology of vertebral degenerative disease in a Pre-Columbian Muisca series from Colombia. *American Journal of Physical Anthropology*, 135: 416–430.
- Ruan D., He Q., Ding Y., Hou L., Li J., and Luk K.D.K. (2007) Intervertebral disc transplantation in the treatment of degenerative spine disease: a preliminary study. *Lancet*, 369: 993–999.
- Sato T., Amano T., Ono H., Ishida H., Kodera H., Matsumura H., Yoneda M., and Masuda R. (2007) Origin and genetic feature of the Okhotsk people, revealed by ancient mitochondrial DNA analysis. *Journal of Human Genetics*, 52: 618–627.
- Sato T., Amano T., Ono H., Ishida H., Kodera H., Matsumura H., Yoneda M., and Masuda R. (2009a) Mitochondrial DNA haplogrouping of the Okhotsk people based on analysis of ancient

- DNA: an intermediate of gene flow from the continental Sakhalin people to the Ainu. *Anthropological Science*, 117: 171–180.
- Sato T., Amano T., Ono H., Ishida H., Koderu H., Matsumura H., Yoneda M., and Masuda R. (2009b) Allele frequencies of the *ABCC11* for earwax phenotypes among ancient populations on the Hokkaido Island, Japan. *Journal of Human Genetics*, 54: 409–413.
- Shedid D. and Benzel E.C. (2007) Cervical spondylosis anatomy: pathophysiology and biomechanics. *Neurosurgery*, 60: S1–S7.
- Shinoda K. (2011) Ancient DNA of the medieval Kamakura people. In: Chujo R., Sakai H., and Ishida H. (eds.), *Natural Science in Archaeology*. Rinsen Shoten, Kyoto, pp. 245–257 (in Japanese).
- Standing S. (ed.) (2008) *Gray's Anatomy. The Anatomical Basis of Clinical Practice*. Churchill Livingstone Elsevier, Edinburgh.
- Suzuki H., Watanabe H., Iwamoto M., Masuda S., Inamoto N., Hayashi T., Tanabe Y., Sakura H., and Kohara Y. (1956) Medieval Japanese skeletons from the burial site at Zaimokuza, Kamakura City. In: *Anthropological Society of Nippon* (ed.), *Zaimokuza Site and Human Skeletons at Kamakura City*. Iwanami Shoten, Tokyo, pp. 1–74 (in Japanese).
- Suzuki T. (1978) A palaeopathological study of the vertebral columns of the Japanese from Jomon to Edo periods. *Journal of the Anthropological Society of Nippon*, 86: 321–336.
- Suzuki T. (1998) Indicators of stress in prehistoric Jomon skeletal remains in Japan. *Anthropological Science*, 106 (Suppl.): 127–137.
- Wada H. (1975) Age-related changes of vertebral body in modern Japanese population. *Sapporo Medical Journal*, 44: 139–152.
- White T.D. (2000) *Human Osteology*, 2nd edn. Academic Press, San Diego.
- Yamaguchi B. (1995) The pectoral and pelvic girdles of the people of the Okhotsk culture from the Omisaki site in Hokkaido. *Bulletin of the National Science Museum, Tokyo, Series D*, 20: 37–46 (in Japanese with English summary).
- Yamaura K. and Ushiro H. (1999) Prehistoric Hokkaido and Ainu origins. In: Fitzhugh W.W. and Dubreuil C.O. (eds.), *Ainu: Spirit of a Northern People*. National Museum of Natural History, Smithsonian Institution, Washington DC, pp. 39–46.
- Yoneda M. (2002) Bones tell about ancient diet. In: Nishiaki Y. and Utagawa H. (eds.), *Northern Aboriginal World*. The University of Tokyo Press, Tokyo, pp. 94–96 (in Japanese).
- Yoneda M., Shibata Y., and Doi N. (2004) Palaeodiet on the Ryukyu Islands: isotope analyses of prehistoric Gusuku and recent periods. *Anthropological Science*, 112: 290.
- Zukeran C., Fukumine T., Doi N., Sensui N., Ishida H., Kanaya F., and Shimabukuro A. (2002) Preliminary observations of some paleopathological conditions in historic and modern human skeletal remains from Ishigaki Island, Ryukyu Islands, Japan. *Anthropological Science*, 110: 421–436.

Appendix 1. Frequencies of osteophytes in anterior and posterior sides of superior and inferior vertebral body surfaces: Okhotsk culture people

		Female						Male					
		Anterior		Differ- ence	Posterior		Sex difference		Anterior		Differ- ence	Posterior	
		F	A/O		F	A/O	Anterior	Posterior	F	A/O		F	A/O
C2	Inferior	31.6	6/19		20.0	4/20			26.7	4/15		40.0	6/15
C3	Superior	11.1	2/18		5.3	1/19			15.4	2/13		7.7	1/13
	Inferior	26.3	5/19		5.0	1/20	<		50.0	7/14		33.3	5/15
C4	Superior	33.3	6/18		10.0	2/20			35.7	5/14		25.0	4/16
	Inferior	63.2	12/19	>>	9.5	2/21	<		41.2	7/17		41.2	7/17
C5	Superior	47.4	9/19	>>	4.8	1/21			36.8	7/19	>	9.5	2/21
	Inferior	68.4	13/19	>>	19.0	4/21			42.1	8/19	>	9.5	2/21
C6	Superior	38.1	8/21		13.0	3/23			42.1	8/19		16.7	4/24
	Inferior	63.2	12/19	>>	9.1	2/22			50.0	10/20		26.1	6/23
C7	Superior	22.7	5/22		7.4	2/27			11.1	2/18		13.0	3/23
	Inferior	26.3	5/19	>	0.0	0/23	>		0.0	0/18		0.0	0/20
T1	Superior	0.0	0/13		0.0	0/19			6.7	1/15		4.8	1/21
	Inferior	0.0	0/16		5.3	1/19			13.6	3/22		12.5	3/24
T2	Superior	6.3	1/16		0.0	0/19			6.3	1/16		0.0	0/21
	Inferior	22.2	4/18		5.3	1/19			15.0	3/20		19.0	4/21
T3	Superior	11.1	2/18		0.0	0/24			18.8	3/16		0.0	0/21
	Inferior	22.2	4/18		4.8	1/21			18.2	4/22		13.0	3/23
T4	Superior	16.7	2/12		0.0	0/18			0.0	0/17		0.0	0/21
	Inferior	33.3	5/15		6.3	1/16			23.8	5/21		26.1	6/23
T5	Superior	17.6	3/17		0.0	0/22			40.0	8/20	>>	0.0	0/22
	Inferior	21.1	4/19	>	0.0	0/20			50.0	11/22	>>	0.0	0/23
T6	Superior	21.1	4/19	>	0.0	0/23			44.4	8/18	>>	0.0	0/23
	Inferior	42.1	8/19	>>	0.0	0/20			40.0	8/20	>>	0.0	0/24
T7	Superior	26.3	5/19	>	0.0	0/25			47.1	8/17	>>	0.0	0/24
	Inferior	40.9	9/22	>>	0.0	0/24			45.0	9/20	>>	0.0	0/23
T8	Superior	38.1	8/21	>>	0.0	0/24			47.1	8/17	>>	0.0	0/18
	Inferior	47.1	8/17	>>	0.0	0/21			41.2	7/17	>>	0.0	0/20
T9	Superior	36.8	7/19	>>	0.0	0/23			38.9	7/18	>>	0.0	0/22
	Inferior	33.3	5/15	>>	0.0	0/20			22.2	4/18	>	0.0	0/23
T10	Superior	31.6	6/19	>>	0.0	0/21			31.6	6/19	>>	0.0	0/26
	Inferior	27.8	5/18		9.5	2/21			20.0	4/20	>	0.0	0/27
T11	Superior	12.5	2/16		0.0	0/21			33.3	8/24	>>	0.0	0/30
	Inferior	27.8	5/18	>	0.0	0/20			17.9	5/28	>	0.0	0/30
T12	Superior	10.0	2/20		0.0	0/22			25.0	4/16	>	0.0	0/24
	Inferior	10.5	2/19		0.0	0/20			18.2	4/22	>	0.0	0/26
L1	Superior	20.0	4/20		0.0	0/19			25.0	4/16	>	0.0	0/20
	Inferior	21.1	4/19		0.0	0/19			27.8	5/18	>	0.0	0/20
L2	Superior	10.5	2/19		0.0	0/20			33.3	6/18	>>	0.0	0/21
	Inferior	27.8	5/18	>	0.0	0/19			39.1	9/23	>>	0.0	0/23
L3	Superior	25.0	4/16	>	0.0	0/20			52.9	9/17	>>	0.0	0/18
	Inferior	37.5	6/16	>>	0.0	0/18			33.3	7/21	>>	0.0	0/22
L4	Superior	33.3	5/15	>>	0.0	0/20	<		73.7	14/19	>>	0.0	0/23
	Inferior	31.6	6/19	>>	0.0	0/20			41.7	10/24	>>	0.0	0/26
L5	Superior	38.9	7/18	>	8.3	2/24			42.9	9/21	>>	4.3	1/23
	Inferior	33.3	7/21		20.8	5/24			9.5	2/21		24.0	6/25
S1	Superior	5.0	1/20		0.0	0/23			18.8	3/16		0.0	0/23

A: number of vertebrae showing lesion (affected), O: number of vertebrae actually observed (observed), F: frequency (%).
 <<, >>, <, > Significantly different at the 0.01 or 0.05 level.

Appendix 2. Frequencies of osteophytes in anterior and posterior sides of superior and inferior vertebral body surface: medieval Kamakura people

		Female						Male					
		Anterior		Differ- ence	Posterior		Sex difference		Anterior		Differ- ence	Posterior	
		F	A/O		F	A/O	Anterior	Posterior	F	A/O		F	A/O
C2	Inferior	1.7	1/58	<	32.2	19/59			3.7	2/54	<<	30.9	17/55
C3	Superior	12.1	7/58		10.5	6/57			8.9	5/57		12.3	7/57
	Inferior	7.1	4/56	<<	25.9	15/58			15.8	9/57	<<	38.6	22/57
C4	Superior	15.0	9/60		10.0	6/60			17.5	10/57		15.8	9/57
	Inferior	11.7	7/60	<	28.3	17/60			10.5	6/57	<	28.1	16/57
C5	Superior	22.8	13/57	>	8.8	5/57			21.8	12/55	>	9.1	5/55
	Inferior	28.6	16/56		28.1	16/57			29.1	16/55		23.6	13/55
C6	Superior	37.5	21/56	>>	12.1	7/58			33.3	19/57	>>	7.0	4/57
	Inferior	27.6	16/58		17.2	10/58			21.1	12/57		14.0	8/57
C7	Superior	23.7	14/59		11.7	7/60			19.0	11/58		8.3	5/60
	Inferior	8.5	5/59		18.6	11/59			1.7	1/59	<<	15.0	9/60
T1	Superior	7.3	4/55		8.9	5/56			9.8	5/51		1.8	1/56
	Inferior	13.0	7/54		20.4	11/54			7.4	4/54		18.2	10/55
T2	Superior	7.7	4/52		3.8	2/53			9.6	5/52		7.0	4/57
	Inferior	11.5	6/52		13.2	7/53			5.9	3/51		16.1	9/56
T3	Superior	8.2	4/49		5.9	3/51			17.4	8/46		9.3	5/54
	Inferior	21.6	11/51	>>	3.8	2/52		<	20.0	10/50		16.4	9/55
T4	Superior	26.0	13/50	>>	1.9	1/52			18.4	9/49	>>	1.8	1/57
	Inferior	25.5	13/51	>	7.5	4/53			26.4	14/53		12.5	7/56
T5	Superior	26.5	13/49	>>	0.0	0/54			30.1	16/53	>>	5.4	3/56
	Inferior	27.5	14/51	>>	1.9	1/54			30.8	16/52	>>	6.9	4/58
T6	Superior	30.2	16/53	>>	0.0	0/55			32.0	17/53	>>	3.5	2/57
	Inferior	37.0	20/54	>>	0.0	0/53			40.0	22/55	>>	6.9	4/58
T7	Superior	41.5	22/53	>>	1.8	1/57			44.6	25/56	>>	1.7	1/59
	Inferior	39.3	22/56	>>	5.4	3/56			43.9	25/57	>>	3.3	2/60
T8	Superior	34.5	19/55	>>	1.8	1/56			49.2	29/59	>>	1.7	1/59
	Inferior	42.6	23/54	>>	5.5	3/55			49.2	29/59	>>	1.7	1/59
T9	Superior	40.7	22/54	>>	0.0	0/54			45.9	28/61	>>	3.3	2/61
	Inferior	46.3	25/54	>>	3.7	2/54			43.5	27/62	>>	3.2	2/62
T10	Superior	36.5	19/52	>>	0.0	0/55			45.2	28/62	>>	0.0	0/64
	Inferior	26.4	14/53	>>	0.0	0/54			32.8	20/61	>>	3.1	2/64
T11	Superior	33.3	19/57	>>	0.0	0/55			39.7	23/58	>>	1.7	1/60
	Inferior	22.8	13/57	>>	1.8	1/57			33.9	20/59	>>	4.8	3/62
T12	Superior	33.9	19/56	>>	0.0	0/57			40	22/55	>>	0.0	0/56
	Inferior	5.4	3/56		1.8	1/56			16.7	10/60	>>	0.0	0/61
L1	Superior	23.8	15/63	>>	0.0	0/61		>	10.2	6/59		1.7	1/59
	Inferior	21.0	13/62	>>	0.0	0/61			11.9	7/59		6.6	4/61
L2	Superior	31.3	20/64	>>	0.0	0/63			29.5	18/61	>>	1.6	1/62
	Inferior	28.3	17/60	>>	0.0	0/60		<	18.0	11/61		9.7	6/62
L3	Superior	51.7	30/58	>>	0.0	0/63			40.7	24/59	>>	4.8	3/63
	Inferior	30.5	18/59	>>	6.5	4/62			21.7	13/60	>>	3.2	2/62
L4	Superior	48.3	29/60	>>	0.0	0/64			42.6	26/61	>>	1.6	1/62
	Inferior	32.8	20/61	>>	4.8	3/62			19.7	12/61		10.0	6/60
L5	Superior	45.6	26/57	>>	8.2	5/61			50	30/60	>>	3.3	2/60
	Inferior	25.4	15/59		13.3	8/60			16.1	10/62		17.7	11/62
S1	Superior	18.2	10/55		8.9	5/56			20	12/60	>>	3.3	2/60

A: number of vertebrae showing lesion (affected), O: number of vertebrae actually observed (observed), F: frequency (%).
 <<, >>, <, > Significantly different at the 0.01 or 0.05 level.

Appendix 3. Frequencies of osteophytes in right and left sides of superior and inferior vertebral body surfaces: Okhotsk culture people

		Female						Male					
		Right		Differ- ence	Left		Sex difference		Right		Differ- ence	Left	
		F	A/O		F	A/O	Right	Left	F	A/O		F	A/O
C2	Inferior	4.8	1/21		9.5	2/21			26.7	4/15		6.3	1/16
C3	Superior	5.3	1/19		20.0	4/20			23.1	3/13		28.6	4/14
	Inferior	5.3	1/19		5.0	1/20			13.3	2/15		16.7	2/12
C4	Superior	15.8	3/19		23.8	5/21			20.0	3/15		41.2	7/17
	Inferior	5.0	1/20		14.3	3/21			0.0	0/17		13.3	2/15
C5	Superior	0.0	0/19		9.5	2/21			10.0	2/20		10.0	2/20
	Inferior	9.5	2/21		13.6	3/22			20.0	4/20		5.3	1/19
C6	Superior	13.6	3/22		18.2	4/22			9.1	2/22		25.0	5/20
	Inferior	9.5	2/21		4.5	1/22			30.0	6/20		13.6	3/22
C7	Superior	4.3	1/23		0.0	0/21		<	20.0	4/20		25.0	5/20
	Inferior	0.0	0/22		0.0	0/23			0.0	0/19		0.0	0/19
T1	Superior	0.0	0/15		0.0	0/17			0.0	0/18		0.0	0/23
	Inferior	0.0	0/17		0.0	0/18			9.5	2/21		0.0	0/19
T2	Superior	0.0	0/16		0.0	0/16			0.0	0/19		0.0	0/20
	Inferior	0.0	0/19		5.3	1/19			0.0	0/20		0.0	0/19
T3	Superior	0.0	0/22		0.0	0/20			5.9	1/17		9.5	2/21
	Inferior	0.0	0/22		4.5	1/22			9.1	2/22		0.0	0/22
T4	Superior	0.0	0/15		0.0	0/13			5.9	1/17		0.0	0/21
	Inferior	12.5	2/16		11.8	2/17			0.0	0/20		0.0	0/18
T5	Superior	0.0	0/19		0.0	0/19			10.5	2/119		0.0	0/22
	Inferior	5.6	1/18		0.0	0/19			13.0	3/23		5.3	1/19
T6	Superior	5.0	1/20		0.0	0/20			9.5	2/21		0.0	0/21
	Inferior	9.1	2/22		4.3	1/23			23.8	5/21		13.6	3/22
T7	Superior	0.0	0/24		0.0	0/22			0.0	0/21		8.7	2/23
	Inferior	7.7	2/26		4.0	1/25			21.7	5/23		13.0	3/23
T8	Superior	0.0	0/22		5.3	1/19			15.8	3/19		0.0	0/19
	Inferior	9.1	2/22		8.7	2/23			15.8	3/19		11.8	2/17
T9	Superior	9.5	2/21		16.7	3/18			0.0	0/17		22.2	4/18
	Inferior	15.0	3/20		14.3	3/21			15.8	3/19		10.0	2/20
T10	Superior	23.8	5/21		22.2	4/18		>	0.0	0/21		0.0	0/23
	Inferior	19.0	4/21		18.2	4/22		>	16.0	4/25		9.5	2/21
T11	Superior	0.0	0/19		5.3	1/19			15.4	4/26		19.2	5/26
	Inferior	9.5	2/21		4.8	1/21			18.5	5/27		6.9	2/29
T12	Superior	5.0	1/20		10.5	2/19			5.3	1/19		21.7	5/23
	Inferior	0.0	0/20		5.0	1/20		>>	32.0	8/25		>	5.6
L1	Superior	26.3	5/19		35.3	6/17			16.7	3/18		15.8	3/19
	Inferior	19.0	4/21		14.3	3/21			22.2	4/18		18.8	3/16
L2	Superior	0.0	0/17		0.0	0/18		<	22.2	4/18		22.7	5/22
	Inferior	10.5	2/19		15.8	3/19			31.8	7/22		15.8	3/19
L3	Superior	6.3	1/16		17.6	3/17		>	38.9	7/18		41.2	7/17
	Inferior	5.6	1/18		11.1	2/18			28.6	6/21		27.8	5/18
L4	Superior	30.0	6/20		38.9	7/18			52.4	11/21		52.2	12/23
	Inferior	10.0	2/20		10.5	2/19			25.0	6/24		18.2	4/22
L5	Superior	38.1	8/21		47.6	10/21			36.4	8/22		33.3	8/24
	Inferior	26.1	6/23		24.0	6/25			39.1	9/23		33.3	7/21
S1	Superior	14.3	3/21		27.3	6/22			13.6	3/22		17.4	4/23

A: number of vertebrae showing lesion (affected), O: number of vertebrae actually observed (observed), F: frequency (%).

<<, >>, <, > Significantly different at the 0.01 or 0.05 level.

Appendix 4. Frequencies of osteophytes in right and left sides of superior and inferior vertebral body surface: medieval Kamakura people

		Female						Male					
		Right		Differ- ence	Left		Sex difference		Right		Differ- ence	Left	
		F	A/O		F	A/O	Right	Left	F	A/O		F	A/O
C2	Inferior	6.9	4/58		8.5	5/59			5.5	3/55		10.9	6/55
C3	Superior	8.8	5/57		8.9	5/56			7.1	4/56		8.9	5/56
	Inferior	5.1	3/59		5.3	3/57	<<	<	21.4	12/56		20.0	11/55
C4	Superior	30.0	18/60		32.7	18/55			23.2	13/56		22.8	13/57
	Inferior	13.3	8/60		10.0	6/60			3.5	2/57		8.9	5/56
C5	Superior	24.6	14/57		21.8	12/55			17.0	9/53		10.9	6/55
	Inferior	17.5	10/57		19.6	11/56			14.5	8/55		16.7	9/54
C6	Superior	31.0	18/58		32.1	18/56		>	20.0	11/55		15.8	9/57
	Inferior	19.0	11/58		22.8	13/57			10.7	6/66		12.5	7/56
C7	Superior	18.6	11/59		17.9	10/56			12.5	7/56		12.5	7/56
	Inferior	10.3	6/58		10.3	6/58			6.8	4/59		3.4	2/59
T1	Superior	1.8	1/57		1.8	1/55			1.8	1/55		1.9	1/54
	Inferior	3.8	2/53		3.8	2/52			7.7	4/52		7.7	4/52
T2	Superior	1.9	1/52		5.9	3/51			3.9	2/51		1.8	1/55
	Inferior	2.0	1/49		1.9	1/53			4.1	2/49		4.0	2/50
T3	Superior	4.0	2/50		0.0	0/49			2.0	1/51		2.0	1/51
	Inferior	7.7	4/52		2.1	1/47			7.7	4/52		2.0	1/51
T4	Superior	8.0	4/50		0.0	0/51			0.0	0/53		0.0	0/55
	Inferior	13.5	7/52		5.9	3/51			7.1	4/56		5.7	3/53
T5	Superior	7.8	4/51		6.0	3/50			7.4	4/54		3.6	2/55
	Inferior	13.5	7/52		3.8	2/52			10.9	6/55		5.6	3/54
T6	Superior	15.4	8/52	>	2.0	1/50			9.8	5/51		5.4	3/56
	Inferior	11.3	6/53		9.8	4/51			3.7	2/54		7.3	4/55
T7	Superior	17.3	9/52	>	3.7	2/54			7.0	4/57		0.0	0/56
	Inferior	16.4	9/55	>>	12.7	1/55			8.6	5/58		8.6	5/58
T8	Superior	11.3	6/53		3.7	2/54			5.2	3/58		1.7	1/58
	Inferior	16.7	9/54		12.7	7/55			10.0	6/60		3.4	2/59
T9	Superior	23.1	12/52		11.3	6/53			6.6	9/61		6.6	4/61
	Inferior	24.1	13/54	>	9.3	5/54			17.7	11/62		13.1	8/61
T10	Superior	21.6	11/51		15.7	8/51			17.5	11/63		8.5	5/59
	Inferior	7.3	4/55		7.7	4/52			18.8	12/64		19.0	12/63
T11	Superior	17.5	10/57		10.7	6/56			19.0	11/58		14.8	9/61
	Inferior	14.0	8/57		14.0	8/57			22.6	14/62		18.3	11/60
T12	Superior	18.2	10/55		10.7	6/56			11.9	7/59		10.3	6/58
	Inferior	5.4	3/56		7.3	4/55	<		18.6	11/59		13.6	8/59
L1	Superior	14.8	9/61		12.7	7/55			4.8	3/62		3.3	2/60
	Inferior	9.8	6/61		11.9	7/59			14.8	9/61		11.9	7/59
L2	Superior	19.4	12/62		25.4	15/59			23.3	14/60		20.6	13/63
	Inferior	21.3	13/61		20.3	12/59			19.4	12/62		17.7	11/62
L3	Superior	40.0	24/60		33.9	20/59			26.2	16/61		31.1	19/61
	Inferior	23.0	14/61		17.7	11/62			14.3	9/63		17.7	11/62
L4	Superior	43.3	26/60		40.0	24/60			31.1	19/61		34.4	21/61
	Inferior	23.0	14/61		26.2	16/61			19.7	12/61		21.7	13/60
L5	Superior	42.4	25/59		41.7	25/60			30.0	18/60		30.0	18/60
	Inferior	27.1	16/59		24.6	15/61		>	14.8	9/61		11.5	7/61
S1	Superior	32.7	18/55		28.3	15/53			23.0	14/61		16.7	10/60

A: number of vertebrae showing lesion (affected), O: number of vertebrae actually observed (observed), F: frequency (%).
 <<, >>, <, > Significantly different at the 0.01 or 0.05 level.

Appendix 5. Frequencies of degenerative changes in right and left sides of superior and inferior apophyseal joints: Okhotsk culture people

		Female						Male					
		Right		Differ- ence	Left		Sex difference		Right		Differ- ence	Left	
		F	A/O		F	A/O	Right	Left	F	A/O		F	A/O
C1	Superior	0.0	0/22		0.0	0/22			11.1	2/18		11.8	2/17
	Inferior	14.3	3/21		4.5	1/22			5.3	1/19		5.9	1/17
C2	Superior	4.3	1/23		4.3	1/23		<	0.0	0/16	<	33.3	6/18
	Inferior	4.3	1/23		4.5	1/22		<	18.8	3/16		31.3	5/16
C3	Superior	13.6	3/22		9.5	2/21			7.1	1/14		14.3	2/14
	Inferior	4.5	1/22		10.0	2/20			9.1	1/11		27.3	3/11
C4	Superior	9.1	2/22		10.0	2/20		<	41.2	7/17		29.4	5/17
	Inferior	9.1	2/22		0.0	0/20			21.4	3/14		18.8	3/16
C5	Superior	4.5	1/22		4.2	1/24		<<	21.1	4/19		41.2	7/17
	Inferior	9.1	2/22		8.7	2/23			12.5	2/16		20.0	3/15
C6	Superior	8.7	2/23		4.3	1/23			21.1	4/19		10.5	2/19
	Inferior	4.3	1/23		0.0	0/24			10.0	2/20		10.0	2/20
C7	Superior	3.6	1/28		0.0	0/27			9.1	2/22		15.0	3/20
	Inferior	3.8	1/26		14.8	4/27			18.2	4/22		21.1	4/19
T1	Superior	13.0	3/23		0.0	0/22		<	29.2	7/24		26.1	6/23
	Inferior	16.7	4/24		4.2	1/24			20.8	5/24		9.1	2/22
T2	Superior	17.4	4/23		13.6	3/22			20.8	5/24		25.0	6/24
	Inferior	18.2	4/22		9.5	2/21			13.0	3/23		4.3	1/23
T3	Superior	16.7	4/24		0.0	0/24		<	25.0	6/24		20.0	5/25
	Inferior	34.8	8/23		22.7	5/22			22.7	5/22		23.8	5/21
T4	Superior	25.0	5/20		10.5	2/19			23.8	5/21		18.2	4/22
	Inferior	44.4	8/18		27.8	5/18			27.8	5/18		27.8	5/18
T5	Superior	33.3	7/21		9.1	2/22			23.8	5/21		28.6	6/21
	Inferior	30.0	6/20		16.7	3/18			21.1	4/19		36.4	8/22
T6	Superior	14.8	4/27		8.3	2/24			13.6	3/22		14.3	3/21
	Inferior	20.0	5/25		9.1	2/22			11.1	2/18		10.5	2/19
T7	Superior	14.8	4/27		11.5	3/26			9.1	2/22		18.2	4/22
	Inferior	7.1	2/28		12.0	3/25			26.3	5/19		22.2	4/18
T8	Superior	20.8	5/24		20.8	5/24			23.8	5/21		15.0	3/20
	Inferior	17.4	4/23		22.7	5/22			20.0	4/20		11.1	2/18
T9	Superior	21.7	5/23		13.0	3/23		<	39.1	9/23		39.1	9/23
	Inferior	28.6	6/21		28.6	6/21			21.1	4/19		36.8	7/19
T10	Superior	9.1	2/22		18.2	4/22			19.2	5/26		25.9	7/27
	Inferior	10.0	2/20		10.0	2/20		<	42.9	9/21	<	36.4	8/22
T11	Superior	21.7	5/23		18.2	4/22			38.5	10/26		38.5	10/26
	Inferior	14.3	3/21		10.0	2/20			20.0	5/25		20.0	5/25
T12	Superior	4.3	1/23		12.5	3/24		<	22.7	5/22	<	37.5	9/24
	Inferior	13.6	3/22		14.3	3/21		<	47.4	9/19	<	50.0	10/20
L1	Superior	22.7	5/22		12.5	3/24		<	57.9	11/19	<	45.0	9/20
	Inferior	17.4	4/23		13.6	3/22		<	16.7	3/18	<	15.8	3/19
L2	Superior	16.7	4/24		20.8	5/24			26.1	6/23		30.4	7/23
	Inferior	20.8	5/24		30.8	8/26			28.6	6/21		19.0	4/21
L3	Superior	17.4	4/23		9.1	2/22		<	45.5	10/22	<	39.1	9/23
	Inferior	18.2	4/22		30.4	7/23		<	28.6	6/21	<	25.0	5/20
L4	Superior	25.9	7/27		19.0	4/21			31.8	7/22		28.0	7/25
	Inferior	42.3	1/261		34.8	8/23			39.1	9/23		47.6	10/21
L5	Superior	29.6	8/27		35.7	10/28			50.0	12/24		47.8	11/23
	Inferior	37.0	1/27		34.6	9/26			35.3	6/17		29.4	5/17
S1	Superior	17.4	4/23		14.3	3/22			33.3	6/18		38.9	7/18

A: number of vertebrae showing lesion (affected), O: number of vertebrae actually observed (observed), F: frequency (%).

<<, >>, <, > Significantly different at the 0.01 or 0.05 level.

Appendix 6. Frequencies of degenerative changes in right and left sides of superior and inferior apophyseal joints: medieval Kamakura people

		Female						Male					
		Right		Differ- ence	Left		Sex difference		Right		Differ- ence	Left	
		F	A/O		F	A/O	Right	Left	F	A/O		F	A/O
C1	Superior	5.5	3/55		1.8	1/56			2.1	1/48		1.9	1/52
	Inferior	7.5	4/53		1.8	1/57			3.9	2/51		3.8	2/53
C2	Superior	3.5	2/57		1.8	1/56			1.8	1/56		0.0	0/55
	Inferior	16.4	9/55		7.0	4/57			7.7	4/52		13.0	7/54
C3	Superior	3.7	2/54		2.0	1/51			7.3	4/55		9.4	5/53
	Inferior	17.3	9/52		8.0	4/50			9.1	5/55		3.7	2/54
C4	Superior	7.1	4/56		8.8	5/57			8.8	5/57		8.9	5/56
	Inferior	15.1	8/53		8.8	5/57			14.0	8/57		5.4	3/56
C5	Superior	3.6	2/56		7.4	4/54			5.9	3/51		3.9	2/51
	Inferior	5.7	3/53		3.6	2/55			0.0	0/52		2.0	1/51
C6	Superior	1.8	1/55		3.6	2/55			5.8	3/52		7.5	4/53
	Inferior	0.0	0/54		0.0	0/54			3.8	2/52		3.8	2/53
C7	Superior	1.8	1/56		0.0	0/55			7.7	4/52		7.8	4/51
	Inferior	3.8	2/53		7.3	4/55			0.0	0/53		1.9	1/52
T1	Superior	8.9	5/56		7.5	4/53			10.7	6/56		7.5	4/53
	Inferior	3.6	2/55		1.8	1/55			1.8	1/56		0.0	0/55
T2	Superior	0.0	0/55		3.8	2/52			0.0	0/52		0.0	0/52
	Inferior	9.8	5/51	>	0.0	0/51			11.5	6/52		7.7	4/52
T3	Superior	2.0	1/50		0.0	0/48			5.9	3/51		4.1	2/49
	Inferior	8.5	4/47		4.1	2/49			10.0	5/50		9.8	5/51
T4	Superior	6.7	3/45		4.3	2/46			15.4	8/52		14.0	7/50
	Inferior	4.3	2/46		2.2	1/45			15.4	8/52		7.7	4/52
T5	Superior	10.4	5/48		4.3	2/46			15.1	8/53		9.8	5/51
	Inferior	10.4	5/48		6.4	3/47			11.8	6/51		9.8	5/51
T6	Superior	2.2	1/46		2.2	1/45	<		13.5	7/52		10.9	5/46
	Inferior	0.0	0/46		2.2	1/45			6.1	3/49		12.2	6/49
T7	Superior	0.0	0/48		0.0	0/46	<<	<<	13.0	7/54		16.4	9/55
	Inferior	10.6	5/47		4.5	2/45			7.5	4/53		7.4	4/54
T8	Superior	2.1	1/48		2.1	1/48	<	<	12.5	7/56		14.8	8/54
	Inferior	13.0	6/46		4.2	2/48			5.4	3/56		10.9	6/55
T9	Superior	6.1	3/49		7.7	4/52			11.1	6/54		12.3	7/57
	Inferior	17.0	8/47		16.0	8/50			13.5	7/52		11.1	6/54
T10	Superior	9.3	5/54		11.1	6/54			12.5	7/56		14.0	8/57
	Inferior	8.0	4/50		5.8	3/52		<	16.7	9/54		17.9	10/56
T11	Superior	11.5	6/52		13.5	7/52			22.2	14/63		27.4	17/62
	Inferior	6.1	3/49		8.0	4/50			13.1	8/61		11.3	7/62
T12	Superior	24.1	13/54		16.4	9/55			22.0	13/59		16.9	10/59
	Inferior	13.7	7/51		9.3	5/54			10.5	6/57		12.1	7/58
L1	Superior	16.4	9/55		7.4	4/54			21.3	13/61		14.5	9/62
	Inferior	18.5	10/54		9.4	5/53			11.3	7/62		9.5	6/63
L2	Superior	7.5	4/53		10.5	6/57			11.5	7/61		8.3	5/60
	Inferior	11.5	6/52		11.1	6/54			10.2	6/59		8.6	5/58
L3	Superior	13.2	7/53		13.0	7/54			10.0	6/60		11.9	7/59
	Inferior	10.7	6/56		8.8	5/57			18.8	12/64		16.1	10/62
L4	Superior	13.8	8/58		10.2	6/59			14.3	9/53		16.7	10/60
	Inferior	25.0	13/52		15.5	9/58			15.0	9/60		13.1	8/61
L5	Superior	12.5	7/56		12.3	7/57			19.7	12/61		11.9	7/59
	Inferior	13.0	7/54		8.8	5/57			10.7	6/56		9.3	5/54
S1	Superior	10.4	5/48		8.9	4/45			5.0	3/60		6.8	4/59

A: number of vertebrae showing lesion (affected), O: number of vertebrae actually observed (observed), F: frequency (%).

<<, >>, <, > Significantly different at the 0.01 or 0.05 level.